

**S.I.S.S.A.**  
**Sector of Functional Analysis and Applications**

*Entrance Examination – October 10<sup>th</sup>, 1994*

Solve at most five of the following problems, explaining your answers.

1. Let  $f \in C^\infty(\mathbf{R})$ .

(i) Prove that for any  $x \in \mathbf{R}$  the Cauchy problem

$$\begin{cases} u' = e^{-f(u)^2} \\ u(0) = x \end{cases}$$

admits a unique solution  $u(\cdot, x)$  defined on  $\mathbf{R}$ .

(ii) Show that  $\lim_{t \rightarrow -\infty} u(t, x) = -\infty$  and  $\lim_{t \rightarrow +\infty} u(t, x) = +\infty$ .

2. Let

$$c_0 = \{(x_n)_{n \in \mathbf{N}} \subset \mathbf{R} : \lim_{n \rightarrow \infty} x_n = 0\}$$

and let  $u : c_0 \rightarrow \mathbf{R}$  be a linear bounded functional. For every  $n \in \mathbf{N}$  set  $\eta_n = u(e^{(n)})$ , where  $e_j^{(n)} = 0$  for  $j \neq n$  and  $e_n^{(n)} = 1$ . Prove that  $(\eta_n) \in l^1$  and  $\|u\| = \sum_{n \in \mathbf{N}} |\eta_n|$ .

3. Let  $f : \mathbf{R} \rightarrow \mathbf{R}$  be a convex function and for every  $x \in \mathbf{R}$  let

$$f'_+(x) = \lim_{h \rightarrow 0^+} \frac{f(x+h) - f(x)}{h}.$$

Show that

$$\int_a^b f'_+(x) ds = f(b) - f(a)$$

for any  $a, b \in \mathbf{R}$  with  $a < b$ .

4. Let  $(X, d)$  be a complete metric space,  $\bar{x} \in X$ ,  $r > 0$  and  $D = \{x \in X : d(x, \bar{x}) \leq r\}$ . Let  $f : D \rightarrow X$  satisfying

$$d(f(x), f(y)) \leq kd(x, y) \quad \text{for any } x, y \in D$$

where  $k \in (0, 1)$  is a constant. Prove that if  $d(\bar{x}, f(\bar{x})) \leq r(1 - k)$  then  $f$  admits a unique fixed point.

5. Let  $I = (0, 1)$  and let  $(f_n)$  be a bounded sequence in  $L^p(I)$  ( $1 < p < \infty$ ). Prove that, if  $(f_n)$  converges to 0 in  $L^1(I)$ , then it converges to 0 in  $L^r(I)$  for any  $r$  with  $1 \leq r < p$ .

6. Characterize the points of  $\mathbf{R}^2$  where the function  $f : \mathbf{R}^2 \rightarrow \mathbf{R}$  defined by

$$f(x, y) = \max\{|x|, |y|\}$$

is differentiable.

7. For any  $\alpha \geq 0$  let  $X^\alpha$  be the space of all functions  $u \in L^2(0, 2\pi)$  whose Fourier coefficients

$$u_n = \int_0^{2\pi} \frac{e^{-int}}{\sqrt{2\pi}} u(t) dt$$

satisfy

$$\sum_{n \in \mathbf{Z}} |n|^{2\alpha} |u_n|^2 < \infty.$$

It is known that  $X^\alpha$  is a Hilbert space with norm

$$\|u\|_\alpha = \left( |u_0|^2 + \sum_{n \in \mathbf{Z}} |n|^{2\alpha} |u_n|^2 \right)^{\frac{1}{2}}.$$

Prove that, if  $0 \leq \alpha < \beta$ , then the imbedding  $X^\beta \hookrightarrow X^\alpha$  is compact.

8. Let  $B = \{(x, y) \in \mathbf{R}^2 : x^2 + y^2 < 1\}$ . Find the solution  $u \in C^2(B) \cap C^0(\bar{B})$  to the problem:

$$\begin{cases} \frac{\partial^2 u}{\partial x^2}(x, y) + \frac{\partial^2 u}{\partial y^2}(x, y) = 0 & \forall (x, y) \in B \\ u(x, y) = 2x^2 & \forall (x, y) \in \partial B. \end{cases}$$

9. Let  $f : \mathbf{R} \rightarrow \mathbf{R}$  be a continuous function. Prove that  $f$  is convex if and only if

$$\int_{-\infty}^{+\infty} f(x)[\varphi(x+h) + \varphi(x-h) - 2\varphi(x)] dx \geq 0$$

for any  $h \in \mathbf{R}$  and for any  $\varphi \in C_0^\infty(\mathbf{R})$  with  $\varphi \geq 0$ .

10. Let  $\mathcal{L}(\mathbf{R}^n, \mathbf{R}^n)$  be the space of all linear maps from  $\mathbf{R}^n$  into  $\mathbf{R}^n$ , let  $A : \mathbf{R}^n \rightarrow \mathcal{L}(\mathbf{R}^n, \mathbf{R}^n)$  be a function of class  $C^1$  and let  $f(x) = A(x)x$  for every  $x \in \mathbf{R}^n$ . Show that if  $A(0)$  is invertible then  $f$  is locally invertible in a neighborhood of 0.